

# AIHA White Paper on Extremely Low Frequency (ELF) Fields

## Introduction

In 1993, The AIHA Board of Directors adopted a position statement on human exposure to extremely low frequency fields. Since that time, members of the AIHA's Nonionizing Radiation Committee have closely followed developments in this area including publication of significant studies of potential health effects and reports of expert panels. Based upon this information, the Nonionizing Radiation Committee has proposed to revise the 1993 Position Statement. The information contained in the white paper is a summary of biological and health effects associated with exposure to ELF fields and present exposure guidelines.

## Physical Characteristics

The ELF spectral region includes the frequencies between 3000 and 3 Hz. In this part of the spectrum the fields are slowly time varying and the wavelengths are quite lengthy. ELF fields are considered as separate, independent, non-radiating electric and magnetic fields at any conceivable observation point.

Electric fields are created by electric charges. The electric field,  $E$ , is defined by the magnitude and direction of the force it exerts on a static unit charge. The units of electric field strength are volts per meter (V/m) although they are often appropriately scaled in kilovolts per meter (kV/m). The magnetic flux density,  $B$ , characterizes the magnetic field strength as  $E$  characterizes the electric field. The units of  $B$  are the tesla (T) that is usually scaled to microtesla (mT) for workplace evaluations. Another unit of  $B$  that may be used is the gauss (G) which is usually scaled to milligauss (mG). [Note: 1 T = 10,000 G.]

Magnetic fields are created by moving charges, or currents. This applies to all fields, whether they are from magnets, power lines, or the earth. Just as the electric field is defined by the force on a unit charge, the magnetic field is defined by the magnitude and direction of the force exerted on a moving charge or current.

## Interaction with Matter/People

Electric fields interact with humans through the outer surface of the body, inducing fields and currents within the body. Hair vibration or other sensory stimuli may occur in fields greater than 10 kV/m. A safety issue arises from currents induced in metal structures, which may produce shocks when humans contact the structure and provide a path to ground.

Time-varying magnetic fields induce electric fields, which in turn induce currents, in tissue in direct proportion to the magnetic flux density, the frequency of oscillation, and the radius of the current loop. The maximum current density induced in the normal residential environment is of the order of microamperes per square meter ( $\mu\text{A}/\text{m}^2$ ). For electric arc welders it may be of the order of milliamperes per square meter ( $\text{mA}/\text{m}^2$ ) or greater.

With photon energy directly proportional to frequency, it is readily apparent that ELF fields will cause neither ionization nor heating. Because the body is a relatively good conductor, the highest internal field that can be induced by an E-field strength in air is about 1 V/m, which leads to a mass-normalized rate of energy transfer of  $10^{-4}$  watts per kilogram (W/kg), 4 orders of magnitude less than the resting metabolic rate (1 W/kg).<sup>(1)</sup> Pulsed magnetic fields can produce higher internal electric fields, but they are still too small to produce measurable tissue heating. Any interactions of ELF fields in air with humans are thus non-thermal.

## Biological and Health Effects

The search for biological and health effects of EMF exposure has been going on at an accelerating pace for more than three decades. Results have been mixed, and their interpretation has been controversial, with little consensus on biological effects and virtually none on health effects that might arise from fields at the levels found in occupational or general community environments.

The next few paragraphs discuss so-called robust biological effects; that is, effects that are reproducible and generally accepted among researchers. This is followed by a discussion of the more controversial non-robust biological effects, which are found by some investigators but not by others who attempt to replicate their results. The section closes with an overview of the epidemiological studies into possible health effects of EMF.

The previous section described how the effect of fields exerts force on charged entities, creating currents, and how the relations between fields and induced currents are quantified through the conductivity of the medium, for example, tissue. Organizations have recommended exposure limits for ELF-EMF based on the following correspondence between current density and biological effects:(2)

- 1-10 mA/m<sup>2</sup>; minor biological effects have been reported;
- 10-100 mA/m<sup>2</sup>; well established effects occur, including effects on the visual and nervous systems;
- 100-1000 mA/m<sup>2</sup>; stimulation of excitable tissue occurs, causing possible health hazards;
- 1000 mA/m<sup>2</sup> and above; extrasystoles and ventricular fibrillation can occur.

Recalling the approximation,  $J \approx 6 B$  (at 60 Hz), magnetic fields of the order of 1 mT would lead to current densities in the range of 6 mA/m<sup>2</sup>.

Phosphenes. This is the sensation of flickering light within the eye. It appears to be produced by stimulation of retinal tissue and by any number of agents (e.g., pressure, mechanical shock, chemical substances, and sudden fright). If the agent inducing phosphenes is ELF fields, then it is termed electrophosphenes and magnetophosphenes, depending on the field that causes them. Phosphenes occur when the induced current density is of the order of 10 mA/m<sup>2</sup> or more. Studies have demonstrated that the sensitivity to phosphenes is greatest around 20 Hz. In general, B-field intensities must be on the order of 10 mT for the production of magnetophosphenes.

Calcium Efflux. In vitro experiments have examined effects on doubly ionized, radioactive calcium. occurring in windows of field frequency and power. A chemical exposure analogy would be as if a little benzene were harmful but a greater amount were not, and a still greater amount were again harmful.

In the first of these studies,(3) a modest increase in the exchange of calcium-45 with a physiologic bath was observed for sectioned chick brain exposed to a 147-MHz carrier wave amplitude modulated at 16 Hz. Other modulation frequencies did not produce as large an effect, and exposure to just a 16-Hz E field had the opposite effect, decreasing efflux. Later work suggested that the effect depended on the relative orientation and strength of the earth's magnetic field, and that it occurred in "windows" of field frequency and intensity and even temperature of the experimental preparation.(4-7)

Genetic Effects. Genetic toxicology studies have found no reliable effects, but there are scattered positive findings.(8-10)

Reproduction and Development. Studies with test animals have examined a variety of species with exposure to E or B fields. There have been scattered positive findings for exposure to test animals, but no consistent, reproducible observations.(11) For example, in a multigenerational study, female miniature swine were exposed to 60-Hz E fields of 35 kV/m for 20 hr/day. When these and an unexposed control group were bred with unexposed males, the control group had a greater rate of fetal malformations than the exposed group. This result reversed in a second breeding. Breeding of the offspring produced similarly conflicting results.(12) Because of the ambiguities in the swine study, a study using a similar protocol was performed with rats (E field = 0, 10, 65, and 130 kV/m).(13) There were no significant increases in litters with malformations among exposed animals.

Six laboratories in North America and Europe examined the effect of B-fields (1  $\mu$ T and pulsed at 100 Hz) on chicken embryos.(14) Data from two laboratories and pooled data from all laboratories found a significant increase in abnormal embryos in the exposed groups. However, the interaction between the incidence of abnormalities and the laboratory doing the experiment was also significant, that is, the effect of exposure differed significantly among laboratories.

Flux densities of 1 or 0.61 mT at 60 Hz were used in two studies of reproductive and developmental toxicology in rats. There was a significant difference the number of fetuses per litter in the 1-mT group in the first experiment, but this was not observed in the replicate study. There were no significant differences in fetal body weight or the

incidence of malformations among the exposure groups.(15) Other researchers exposed pregnant rats (0.002, 0.2, and 1 mT at 60 Hz) and found no significant differences in developmental abnormalities.(16)

Reviews of human studies are available.(11,17) Evaluations examined residential exposure (appliances esp. electric blankets), occupational exposure other than VDT use (e.g., substations), and VDT use. End points evaluated include spontaneous abortion, congenital malformations, preterm delivery, low birth weight, and intrauterine growth retardation. A few scattered positive findings have been reported, but no trends established. However, there have been relatively few studies of exposures to sources other than to VDT operation. There are a number of methodological limitations which are addressed in the review articles cited above.

Effect on Melatonin. The possible effects of ELF fields on levels of the pineal hormone melatonin have been investigated. Animal studies have demonstrated effects on melatonin from E fields,(18,19 B fields,(20) and no effects.(21-24) Other effects include E-field exposure decreased serum melatonin but not pineal melatonin levels;(25) pulsed static B fields affected melatonin if exposure was during the mid- or late-dark phase, but not when exposure was during the day time or early dark phase;(26) short-term B-field exposure affected pineal and nocturnal serum melatonin, while long-term exposure affected just nocturnal serum melatonin;(27) serum melatonin was not affected by daytime E- and B-field exposure with a slow onset,(28) but it was reduced by variable exposure with a rapid onset/offset.(29) The National Institute of Environmental Health Sciences (NIEHS) Working Group concluded that there is weak evidence that exposure to E and B fields alters melatonin in rodents but not in sheep or baboons.(30)

The outcome of twelve studies on possible ELF-induced effects on melatonin in humans has been reviewed by the NIEHS. Five laboratory studies were negative, but one laboratory and six observational studies demonstrated some effects on melatonin or its urinary metabolite, 6-hydroxymelatonin sulfate (6-OHMS).(30) In later studies, no statistically significant effects were reported in female garment factory workers,(31) men exposed to an electric sheet,(32) and women(33) and men exposed to intense B fields at night.(34) A decrease in nocturnal melatonin was observed in women in another study,(35) while others have suggested a reduction in 6-OHMS associated with work in an electrical substation or with 3-phase fields.(36)

Cancer Studies[1]. A number of cancer types have been evaluated but the types studied most often in relation to ELF exposure are leukemia, brain cancer, and breast cancer. Experiments with test animals have examined initiation, promotion and co-promotion in models of mammary cancer, skin cancer, brain cancer, leukemia, and lymphoma. The findings are complex to interpret and do not demonstrate any clear trends. McCann and colleagues reviewed the published literature and concluded that "a weak promoting effect of Mfs [magnetic fields] under certain exposure conditions cannot be ruled out categorically based on available data."(37) Boorman et al. examined the literature on mammary cancer finding that "The totality of rodent data does not support the hypothesis that 50 or 60 Hz magnetic-field exposure enhances mammary cancer in rodents, nor does it provide experimental support for possible epidemiological associations between magnetic-field exposure and increased breast cancer risk."(38)

[1] Note: Because of the large number and complex nature of published studies, this section will draw largely upon findings from critical reviews published in either the peer-reviewed literature or by expert panels.

For human studies, evaluations have included both residential and occupational settings and children and adults. A number of useful review articles and meta-analyses are available including general(30,39-42) and topical reviews of leukemia,(43-49) brain cancer,(44,50,51) and breast cancer.(52-54)

Early occupational studies, here defined as prior to 1993, often utilized job titles and information from death certificates. In general, many of these studies found no-to-modest increases in risk of leukemia and brain cancer and no dose-response effect.(44)

Early studies of childhood cancer utilized wire coding to classify residences with regards to potential magnetic field intensity, and spot measurements. Generally, these studies demonstrated a weak association with leukemia.(611)

More recent studies have utilized spot and personal measurement strategies and detailed numerical models to estimate exposure. Three studies examined the risk of power company workers in the U.S., Canada, and France (see Table below). Of these, one reported no increase in risk of leukemia and brain cancer,(55) while the others found an increase in risk of leukemia(56) and brain cancer.(57) A combined analysis of the data from the three studies found that the relative risk (RR) per 10 mT-yrs was 1.09 (95% CI = 0.98 to 1.21) for leukemia and 1.12 (95% confidence interval, CI = 0.98 to 1.28) for brain cancer.(45)

Table - Summary of Three Power Industry Studies

Study	Sahl et al. (1993) <sup>(55)</sup>	Theriault et al. (1994) <sup>(56)</sup>	Savitz et al. (1995) <sup>(57)</sup>
Exposure	Above mean	Above 90th (15.7 $\mu$ T-years)	Above 90th measure $\mu$ T-work years or 18.8 $\mu$ T-years
Brain cancer RR (95% CI)	0.81 (0.48-1.36)	1.95 (0.76-5.00)	2.29 (1.15-4.56)
Leukemia RR (95% CI)	1.07 (0.80-1.45)	1.75 (0.77-3.96)	1.11 (0.57-2.14)
AML RR (95% CI)	-----	2.68 (0.50-14.50)	1.62 (0.51-5.12)

A meta-analysis of 38 studies of occupational B-field exposure and leukemia reported a pooled RR = 1.18 (CI = 1.12-1.124). The highest pooled RR (1.55, CI = 1.10-2.19) was for the chronic lymphocytic leukemia subtype (12 studies).(46) The NIEHS Working Group found limited evidence of an increased risk of chronic lymphocytic leukemia in workers and exposure to ELF magnetic fields.(30) The Advisory Group on Non-ionizing Radiation (AGNIR) of the United Kingdom's National Radiological Protection Board (NRPB), concluded that a review of occupational cohort studies on leukemia and ELF exposure are equivocal, at best.(41)

A pooled analysis of childhood leukemia found no increase in risk for B-fields < 0.4 mT, while the RR = 2.00 (95% CI = 1.27 to 3.13) for children with residential exposure  $\geq$ 0.4 mT.(47) Wartenberg performed a meta-analysis for childhood leukemia finding the following RR values: 1.1 (0.9-1.3) for spot measurements, 1.2 (0.9-1.5) for calculated fields, 2.7 (0.8-8.7) for wire codes by spot measurements, and 1.6 (0.5-4.6) for wire codes by 24-hour measurements. He estimates 175 cases of leukemia by wire code data and 240 cases by spot measurements with the following assumptions: exposure causes leukemia; the studies are accurate and representative; and that exposure follows a log-linear relationship.(49) In a review of the literature, both the NIEHS Working Group and the International Agency for Research on Cancer (IARC) found limited evidence for an increased risk of childhood leukemia and residential exposure.(42)

Kheifets and colleagues have reviewed the literature on brain cancer. In meta-analyses she reported a pooled RR = 1.21 (CI = 1.11-1.33) in 1995(50) and a pooled OR = 1.16 (95% CI = 1.08-1.24) in 2001. In the latter review, Kheifets states that there is "little support for an association between residential EMF exposure and childhood or adult brain cancer."(51) In a review of three major studies of utility workers, the estimated combined RR/10 mT-yr was 1.12 (95% CI = 0.98-1.28), as noted above.(45) Members of the NIEHS Working Group(30) and IARC(42) concluded that there was inadequate evidence of brain cancer in adults and children, while AGNIR concluded evidence from cohort studies was equivocal.(41)

The risk of breast cancer and ELF exposure has been evaluated in men and pre-, peri-, and postmenopausal women. Erren reviewed 48 published studies and included studies with estimates of RR (24 for women and 15 for men) in a meta-analysis. The average RR = 1.12 (95% CI = 1.09-1.15) for women and 1.37 (95% CI = 1.11-1.71) for men demonstrates a small increase in risk of breast cancer.(52) After reviewing the literature, other researchers have suggested that this is an area requiring further research.(53,54)

Over the years, a number of expert panels have addressed the question of ELF exposure and cancer. Three of these mentioned earlier are the NIEHS Working Group, IARC, and AGNIR. All three groups examined an extensive volume of scientific literature in their assessments. Based upon their evaluations, the NIEHS Working Group classified ELF magnetic fields as a Group 2B (possibly) carcinogen for childhood leukemia and chronic lymphocytic leukemia in workers.

IARC found limited evidence for an excess cancer risk on the basis of childhood leukemia from exposure to high residential magnetic fields, and also placed ELF magnetic fields in Group 2B. IARC found inadequate evidence for all other cancer types in children and adults from exposure to ELF electric and magnetic fields.(42) AGNIR, led by Sir Richard Doll, found "some epidemiological evidence that prolonged exposure to higher magnetic fields is associated with a small risk of leukemia in children."(41) Both IARC and AGNIR cite a value of  $\geq$ 0.4 mT when referring to high levels of residential exposure.

## Exposure Guidelines

Ideally, exposure guidelines and standards are established on the basis of an accepted mechanism of interaction, dose-response studies in animals, and epidemiological evidence of similar effects in humans. None of this has occurred for ELF fields. No accepted, biologically plausible mechanism has been advanced to explain how fields interact with biological systems to yield observed in vitro responses, much less disease in an organism. In fact, the field parameter or parameters to be measured, because of their possible biological significance, are unknown. The traditional approaches that "more is worse" and that time-weighted-average or even peak exposures are the quantities of interest have been called into question by studies suggesting that effects occur in windows of frequency and power. The choice of exposure metric is thus made difficult by the lack of knowledge about interaction mechanisms and the lack of a clearly defined, exposure-associated health effect.

The current biological basis for the exposure guidelines includes acute effects to central nervous system tissues. Effects that are often cited include phosphenes and alterations of visual evoked potentials (VEP). These effects occur at relatively high levels of exposure. Phosphenes is a biological effect while alterations of the VEP "are not necessarily harmful."(58)

Exposure guidance, based on the current understanding of ELF-induced biological effects, has been developed by a number of countries and organizations.(58,59) This chapter will focus on guidelines published by the ACGIH(60) and the ICNIRP.(61,62)

The rationale for these exposure limits is based on induced body currents. For occupational exposures, both the ACGIH and ICNIRP recommend limiting induced current densities in the body to those levels that occur normally, that is, up to about 10 mA/m<sup>2</sup> (higher current densities can also occur naturally in the heart). They acknowledged that biological effects have been demonstrated in laboratory studies at field strengths below those permitted by the exposure guidelines, but both concluded that there was no convincing evidence that occupational exposure to these field levels leads to adverse health effects.

The ACGIH guideline extends into part of the RF spectral region, which that organization calls subradiofrequency (sub-RF).(60) The ACGIH TLV for occupational exposure to ELF electric fields states that exposure should not exceed 25 kV/m for frequencies from 0 (DC) to 100 Hz. For frequencies in the range of 100 Hz to 4 kHz, the TLV is given by

$$ETLV = 2.5 \times 10^6 / f \text{ (V/m)}$$

where E has units of V/m (rms value) and f is frequency in Hz. A proviso is added for workers with cardiac pacemakers, limiting power-frequency exposures to 1 kV/m. Electromagnetic interference with pacemaker function may occur in some models at power-frequency electric fields as low as 2 kV/m.

The TLV for magnetic fields from 1 to 300 Hz limits routine occupational (rms) exposure to ceiling values determined by

$$BTLV = 60 / f \text{ (mT)}$$

where f is the frequency in Hz. At frequencies below 1 Hz, the TLV is 60 mT. From 300 Hz to 30 kHz the ceiling value is a limit of 0.2 mT. For workers with cardiac pacemakers, the limit is 0.1 mT at power frequencies.

Because of ambiguities in the scientific understanding of the mechanism of biological interaction, the sub-RF TLVs are ceiling limits, that is, they are values not to be exceeded.(60)

ICNIRP makes recommendations for exposure limits of both workers and members of the general public to electric and magnetic fields.(61) The fundamental quantity, the current density, is called the basic restriction and the measured field quantity is the reference level. As mentioned above, the basic restriction for the occupational standard is a current density of 10 mA/m<sup>2</sup>, while the basic restriction for the general public is 2 mA/m<sup>2</sup> for frequencies between 4 Hz and 1 kHz. At frequencies between 1 and 4 Hz, the current density is frequency dependent. ICNIRP does not recommend an averaging time for the ELF spectral region "because the known effects of induced and contact currents at those frequencies are acute phenomena involving a rapid response of the nervous system."(62)

Frequency-dependent reference levels are recommended both E and B. The allowable E-field strength reaches a maximum in the band between 1 and 25 Hz at values of 20 and 10 kV/m for occupational and general public exposure, respectively. Field strength then decreases with frequency before reaching another plateau. Exposure limits at 60 Hz are 8333 V/m for workers and 4167 V/m for members of the general public.

For B fields, the maximum flux density occurs at less than 1 Hz, then decreases in a frequency-dependent manner up to around 800 Hz. Exposure limits at 60 Hz are 417 and 83  $\mu$ T for workers and the general public, respectively.

## Exposure Assessment

Although not addressed in this report, information on evaluating exposures to ELF fields is available in two AIHA publications: The Occupational Environment -- Its Evaluation and Control (63) and the Nonionizing Radiation Guide Series, Extremely Low Frequency (ELF) Electric and Magnetic Fields.(64) In addition to information on instruments and measurements, these two publications include reviews of key issues dealing with ELF exposure. Also, the National Institute for Occupational Safety and Health (NIOSH) has published a useful guide, NIOSH Manual for Measuring Occupational Electric and Magnetic Field Exposures.(65)

Adopted by AIHA Board of Directors – June 15, 1993

Updated and Adopted by AIHA Board of Directors – August 4, 2002

## References

1. Tenforde, T.S.: Biological Interactions of Extremely-Low-Frequency Electric and Magnetic Fields. *Biochem. Bioenergetics* 25:1-17 (1991).
2. Bernhardt, J.H., H.J. Haubrich, G. Newi, N. Krause, and K.H. Schneider: Limits for Electric and Magnetic Fields in DIN VDE Standards: Considerations for the Range 0 to 10 kHz. CIGRE, International Conference on Large High Voltage Electric Systems. 112 Boulevard Haussmann, 75008 Paris, August 27th-September 4th, 1986.
3. Bawin, S.M., L.K. Kaczmarek, and W.R. Adey: Effects of Modulated VHF Fields on the Central Nervous System. *Ann. N.Y. Acad. Sci.* 247:74-81 (1975).
4. Blackman, C.F., S.G. Benane, L.S. Kinney, W.T. Joines, and D.E. House: Effects of ELF Fields on Calcium-Ion Efflux from Brain Tissue in vitro. *Radiat. Res.* 92:510-520 (1982).
5. Blackman, C.F., S.G. Benane, J.R. Rabinowitz, D.E. House, W.T. Joines: A Role for the Magnetic Field in the Radiation-Induced Efflux of Calcium Ions from Brain Tissue in vitro. *Bioelectromagnetics* 6:327-337 (1985).
6. Blackman, C.F., L.S. Kinney, D.E. House, and W.T. Joines: Multiple Power-Density Windows and Their Possible Origin. *Bioelectromagnetics* 10:115-128 (1989).
7. Blackman, C.F., S.G. Benane, D.E. House: The Influence of Temperature During Electric- and Magnetic-Field-Induced Alteration of Calcium-Ion Release from in vitro Brain Tissue. *Bioelectromagnetics* 12:173-182 (1991).
8. Murphy, J.C., D.A. Kaden, J. Warren, and A. Sivak: Power Frequency Electric and Magnetic Fields: A Review of Genetic Toxicology. *Mutation Res.* 296:221-240 (1993).
9. McCann, J., F. Dietrich, C. Rafferty, and A.O. Martin: A Critical Review of the Genotoxic Potential of Electric and Magnetic Fields. *Mutation Res.* 297:61-95 (1993).
10. McCann, J., F. Dietrich, and C. Rafferty: The Genotoxic Potential of Electric and Magnetic Fields: An Update. *Mutation Res.* 411: 45-86 (1998).
11. Huuskonen, H., M.L. Lindbohm, and J. Juutilainen: Teratogenic and Reproductive Effects of Low-Frequency Magnetic-Fields. *Mutation Res.* 410: 167-183 (1998).

12. Sikov, M.R., D.N. Rommereim, J.L. Beamer, R.L. Buschbom, W.T. Kaune, and R.D. Phillips: Developmental Studies of Hanford Miniature Swine Exposed to 60-Hz Electric Fields. *Bioelectromagnetics* 8:229-242 (1987).
13. Rommereim, D.N., R.L. Rommereim, L.E. Anderson, and M.R. Sikov: Reproductive and Teratologic Evaluation in Rats Chronically Exposed at Multiple Strengths of 60-Hz Electric Fields. In Abstracts, Tenth Annual Meeting of the Bioelectromagnetics Society, Stamford, CT, 1988.
14. Berman, E., L. Chacon, D. House, B.A. Koch, W.E. Koch, J. Leal, S. Lovtrup, E. Mantiply, A.H. Martin, G.I. Martucci, K.H. Mild, J.C. Monahan, M. Sandstrom, K. Shamsaifar, R. Tell, M.A. Trillo, A. Ubeda, and P. Wagner: Development of Chicken Embryos in A Pulsed Magnetic Field. *Bioelectromagnetics* 11:169-187 (1990).
15. Rommereim, D.N., R.L. Rommereim, D.L. Miller, R.L. Buschbom, and L.E. Anderson: Developmental Toxicology Evaluation of 60-Hz, Horizontal Magnetic Fields in Rats. *Appl. Occupat. Environm. Hyg.* 11: 307-312 (1996).
16. Ryan, B.M., E. Mallett, T.R. Johnson, J.R. Gauger, and D.L. McCormack: Developmental Toxicity Study of 60 Hz (Power Frequency) Magnetic Fields in Rats. *Teratology* 54: 73-83 (1996).
17. Shaw, G.M.: Adverse Human Reproductive Outcomes and Electromagnetic Fields: A Brief Summary of the Epidemiologic Literature. *Bioelectromagnetics* 5: S5-S18 (2001).
18. Wilson, B.W., L.E. Anderson, D.I. Hilton, and R.D. Phillips: Chronic Exposure to 60-Hz Electric Fields: Effects on Pineal Function in the Rat. *Bioelectromagnetics* 2:371-380 (1981).
19. Wilson, B.W., E.K. Chess, and L.E. Anderson: 60-Hz Electric-Field Effects on Pineal Melatonin Rhythms: Time Course for Onset and Recovery. *Bioelectromagnetics* 7:239-242 (1986).
20. Kato, M., K. Honma, T. Shigemitsu, and Y. Shiga: Effects of Exposure to a Circularly Polarized 50-Hz Magnetic Field on Plasma and Pineal Melatonin Levels in Rats. *Bioelectromagnetics* 14:97-106 (1993).
21. Lee, J.M., F. Stormshak, J.M. Thompson, P. Thinesen, L.J. Painter, E.G. Olenchek, D.L. Hess, R. Forbes, and D.L. Foster: Melatonin Secretion and Puberty in Female Lambs Exposed to Environmental Electric and Magnetic Fields. *Biol. Reprod.* 49:857-864 (1993).
22. Lee, J.M., F. Stormshak, J.M. Thompson, D.L. Hess, and D.L. Foster: Melatonin and Puberty in Female Lambs Exposed to EMF: A Replicate Study. *Bioelectromagnetics* 16:119-123 (1995).
23. Truong, H., and S.M. Yellow: Continuous or Intermittent 60 Hz Magnetic Field Exposure Fails to Affect the Nighttime Rise in Melatonin in the Adult Djungarian Hamster. *Biol. Reprod.* 52:72 (1995).
24. Levine, R.L., J.K. Dooley, and T.D. Bluni: Magnetic-Field Effects on Spatial Discrimination and Melatonin Levels in Mice. *Physiol. Behav.* 58:535-537 (1995).
25. Grota, L.J., R.J. Reiter, P. Keng, and S. Michaelson: Electric Field Exposure Alters Serum Melatonin but Not Pineal Melatonin Synthesis in Male Rats. *Bioelectromagnetics* 15:427-438 (1994).
26. Yaga, K. R.J. Reiter, L.C. Manchester, H. Nieves, J.H. Sun, and L.D. Chen: Pineal Sensitivity to Pulsed Static Magnetic-Fields Changes during the Photoperiod. *Brain Res. Bull.* 30:153-156 (1993).
27. Selmaoui, B., and Y. Touitou: Sinusoidal 50-Hz Magnetic-Fields Depress Rat Pineal NAT Activity and Serum Melatonin. Role of Duration and Intensity of Exposure. *Life Sci.* 57:1351-1358 (1995).
28. Rogers, W.R., R.J. Reiter, L. Barlow-Walden, H.D. Smith, and J.L. Orr: Regularly Scheduled, Day-Time, Slow Onset 60 Hz Electric and Magnetic Field Exposure Does Not Depress Serum Melatonin Concentration in Nonhuman Primates. *Bioelectromagnetics (Suppl.)* 3:111-118 (1995).

29. Rogers, W.R., R.J. Reiter, H.D. Smith, and L. Barlow-Walden: Rapid-Onset/Offset, Variably Scheduled 60 Hz Electric and Magnetic Field Exposure Reduces Nocturnal Serum Melatonin Concentration in Nonhuman Primates. *Bioelectromagnetics (suppl) 3*:119-122 (1995).
30. National Institute of Environmental Health Sciences: Assessment of Health Effects from Exposure to Power-Line Frequency Electric and Magnetic Fields (NIEHS Working Group Report). Research Triangle Park, NC: NIEHS: 1998.
31. Juutilainen, J., R.G. Stevens, L.E. Anderson, N.H. Hansen, M. Kilpelainen, T. Kumlin, J.T. Laitinen, E. Sobel, and B.W. Wilson: Nocturnal 6-hydroxymelatonin sulfate Excretion in Female Workers Exposed to Magnetic Fields. *J. Pineal Res.* 28: 97-104 (2000).
32. Hong, S.C., Y. Kurokawa, M. Kabuto, and R. Ohtsuka: Chronic Exposure to ELF Magnetic Fields During Night Sleep with Electric Sheets: Effects on Diurnal Melatonin Rhythms in Men. *Bioelectromagnetics* 22: 138-143 (2001).
33. Graham, C., M.R. Cook, M.M. Gerkovich, and A. Sastre: Examination of the Melatonin Hypothesis in Women Exposed at Night to EMF or Bright Light. *Environ. Health. Perspect.* 109: 501-507 (2001a).
34. Graham, C., M.R. Cook, M.M. Gerkovich, and A. Sastre: Melatonin and 6-OHMS in high-intensity magnetic fields. *J. Pineal Res.* 31: 85-88 (2001).
35. Davis, S., W.T. Kaune, D.K. Mirick, C. Chen, and R.G. Stevens: Residential Magnetic Fields, Light-at-Night, and Nocturnal Urinary 6-Sulfatoxymelatonin Concentration in Women. *Am. J. Epidemiol.* 154: 591-600 (2001).
36. Burch, J.B., J.S. Reif, C.W. Noonan, and M.G. Yost: Melatonin Metabolite Levels in Workers Exposed to 60-Hz Magnetic Fields: Work in Substations and with 3-Phase Conductors. *J. Occup. Environ. Med.* 42: 136-142 (2000).
37. McCann, J., R. Kavet, and C.N. Rafferty: Assessing the Potential Carcinogenic Activity of Magnetic Fields Using Animal Models. *Environm. Health Perspect.* 108(suppl 1): 79-100 (2000).
38. Boorman, G.A., D.L. McCormick, J.M. Ward, J.K. Haseman, and R.C. Sills: Magnetic Fields and Mammary Cancer in Rodents: A Critical Review and Evaluation of Published Literature. *Radiat. Res.* 153: 617-626 (2000).
39. Heath, C.W.: Electromagnetic Field Exposure and Cancer: A Review of Epidemiologic Evidence. *Ca. Cancer J. Clin.* 65:29-44 (1996).
40. Carstensen, E.L.: Magnetic Fields and Cancer. *IEEE Eng. Med. Biol. Mag.* 14(4):362-369 (1995).
41. Advisory Group on Non-ionising Radiation: ELF Electromagnetic Fields and the Risk of Cancer. Doc. NRPB: 12(1-179) 2001.
42. International Agency for Research on Cancer: Static and Extremely Low Frequency Electric and Magnetic Fields, vol. 80 IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Lyon, France: IARC (in preparation 2002) <http://193.51.164.11/htdocs/announcements/vol80.htm>
43. Feychting, M.: Occupational Exposure to Electromagnetic Fields and Adult Leukemia: A Review of the Epidemiological Evidence. *Radiat. Environ. Biophys.* 35:237-242 (1996).
44. Miller, R.D., J.S. Neuberger, and K.B. Gerald: Brain Cancer and Leukemia and Exposure to Power-Frequency (50-to 60-Hz) Electric and Magnetic Fields. *Epidemiol. Rev.* 19: 273-293 (1997).
45. Kheifets, L.I., E.S. Gilbert, S.S. Sussman, P. Gunel, J.D. Sahl, D. A. Savitz, and G. Theriault: Comparative Analyses of the Studies of Magnetic Field and Cancer in Electric Utility Workers: Studies from France, Canada, and the United States. *Occup. Environ. Med.* 56: 567-574 (1999).
46. Kheifets, L.I., A.A. Afifi, P.A. Buffler, Z.W. Zhang, and C.C. Matkin: Occupational Electric and Magnetic Field Exposure and Leukemia. *J. Occup. Environ. Med.* 39: 1074-1091 (1997).



47. Ahlbom, A., N. Day, M. Feychting, E. Roman, J. Skinner, M. Kinet, M. McBride, J. Michaelis, J.H. Olsen, T. Tynes, and P.K. Verkasalo: A Pooled Analysis of Magnetic Fields and Childhood Leukaemia. *Brit. J. Cancer* 83: 692-698 (2000).
48. Greenland, S., A.R. Sheppard, W.T. Kaune, C. Poole, and M.A. Kelsh: A Pooled Analysis of Magnetic Fields, Wire Codes, and Childhood Leukemia. *Epidemiology* 11: 624-634 (2000).
49. Wartenberg, D.: Residential EMF Exposure and Childhood Leukemia: Meta-Analysis and Population Attributable Risk. *Bioelectromagnetics Supplement 5*: S86-S104 (2001).
50. Kheifets, L.I., A.A. Afifi, P.A. Buffler, and Z.W. Zhang: Occupational Electric and Magnetic Field Exposure and Brain Cancer: A Meta-Analysis. *J. Occupat. Environ. Med.* 37:1327-1341 (1995).
51. Kheifets, L.I.: Electric and Magnetic Field Exposure and Brain Cancer: A Review. *Bioelectromagnetics S5*: S120-S131 (2001).
52. Erren, T.C.: A meta-Analysis of Epidemiologic Studies of Electric and Magnetic Fields and Breast Cancer in Women and Men. *Bioelectromagnetics S5*:S105-S119 (2001).
53. Brainard, G.C., R. Kavet, and L.I. Kheifets: The Relationship between Electromagnetic Field and Light Exposures to Melatonin and Breast Cancer: A Review of the Relevant Literature. *J. Pineal Res.* 26: 65-100 (1999).
54. Caplan, L.S., E.R. Schoenfeld, E.S. O'Leary, and M.C. Leske: Breast Cancer and Electromagnetic Fields--A Review. *Ann. Epidemiol.* 10: 31-44 (2000).
55. Sahl, J.D., M.A. Kelsh, and S. Greenland: Cohort and Nested Case-Control Studies of Hematopoietic Cancers and Brain Cancers among Electric Utility Workers. *Epidemiology* 4:104-14 (1993).
56. Theriault, G., M. Goldberg, A.B. Miller, B. Armstrong, P. Guenel, J. Deadman, E. Imbernon, T. To, A. Chevallier, D. Cyr, and C. Wall: Cancer Risks Associated with Occupational Exposure to Magnetic Fields among Electric Utility Workers in Ontario and Quebec, Canada, and France: 1970-1989. *Am. J. Epidemiol.* 139:550-72 (1994).
57. Savitz, D.A., and D. P. Loomis: Magnetic Field Exposure in Relation to Leukemia and Brain Cancer Mortality among Electric Utility Workers. *Am. J. Epidemiol.* 141:123-34 (1995).
58. Bailey, W.H., S.H. Su, T.D. Bracken, and R. Kavet: Summary and Evaluation of Guidelines for Occupational Exposure to Power Frequency Electric and Magnetic Fields. *Health Phys.* 73: 433-453 (1997).
59. Environment & Society Working Group, Union of the Electricity Industry: Status Report Power-Frequency EMF Exposure Standards (Ref: 2001-2650-0010). Brussels, Belgium: Eurelectric, October 2001.
60. American Conference of Governmental Industrial Hygienists: 2001 TLVs and BEIs. Cincinnati: American Conference of Governmental Industrial Hygienists, 2001.
61. International Commission on Non-Ionizing Radiation Protection: Guidelines for Limiting Exposure to Time-Varying Electric, Magnetic, and Electromagnetic Fields (up to 300 GHz). *Health Phys.* 74: 494-522 (1998).
62. Mathes, R.: Response to Questions and Comments on ICNIRP Guidelines for Limiting Exposure to Time-Varying Electric, Magnetic, and Electromagnetic Fields (up to 300 GHz). *Health Phys.* 75: 438-439 (1998).
63. Hitchcock, R.T., C. E. Moss, W.E. Muiray, R.M. Patterson, and J.R. Rockwell, Jr.: Nonionizing Radiation. In *The Occupational Environment -- Its Evaluation and Control*, 2nd Ed. Fairfax, VA: American Industrial Hygiene Association, 2002.
64. Hitchcock, R.T., S. McMahan, and G.C. Miller: Extremely Low Frequency (ELF) Electric and Magnetic Fields. Fairfax, VA: American Industrial Hygiene Association, 1995.

65. National Institute for Occupational Safety and Health: Manual for Measuring Occupational Electric and Magnetic Field Exposures (DHHS (NIOSH) Publication No. 98-154) by J.D. Bowman, M.A. Kelsh, and W.T. Kaune. Cincinnati: NIOSH, 1998.